



SUMMER CAMP

Steps that summer camps can take to reduce the risk of spreading COVID-19.

- Promote, and consider requiring that all staff and eligible campers get vaccinated. Getting a vaccination is the best way to protect yourself, prevent the spread of COVID-19, and protect others.
- Ask unvaccinated campers and staff members to engage in a 2-week prearrival quarantine that includes physical distancing, mask wearing when not at home, avoiding unnecessary travel, and refraining from indoor social gatherings with people outside of their households.
- Ask campers and staff who are not fully vaccinated to provide proof of a negative viral test taken no more than 1–3 days before their arrival at camp. Delay arrival for campers or staff with confirmed positive test results.
- Assign campers to cohorts or groups.
- Require unvaccinated individuals to wear face coverings indoors.
- Immediately isolate sick campers and staff away from others.
- Clean surfaces once a day, prioritizing high-touch surfaces.

More actions camps can take...

- Teach and reinforce hand hygiene for adults and campers such as washing hands frequently with soap and water for at least 20 seconds (about as long as it takes to sing “Happy Birthday” twice).
- Perform activities outdoors that have the potential to produce respiratory droplets, including singing, chanting, shouting, or playing an instrument.
- Stagger meal service so that physical distancing between different cohorts can be maintained.

