



NC State Health Director’s Statewide Standing Order
Administration of Moderna COVID-19 Vaccine Bivalent Original and Omicron BA.4 /BA. 5 in Individuals 18 Years of Age and Older
September 2, 2022

Purpose: To reduce morbidity and mortality from coronavirus disease 2019 (COVID-19) by vaccinating persons who meet the criteria established by the Centers for Disease Control and Prevention’s Advisory Committee on Immunization Practices (ACIP) and to administer Moderna COVID-19 Vaccine, Bivalent (Original and Omicron BA.4/BA.5), herein referred to Moderna COVID-19 Vaccine, Bivalent.

NOTE: On September 1, 2022, CDC recommended the use of Moderna COVID-19 Vaccine Bivalent Original and Omicron BA.4/BA.5 as authorized by FDA EUA on August 31, 2022.

Policy: This standing order authorizes any North Carolina healthcare provider, in accordance with the conditions of their licensure, or pursuant to orders issued under North Carolina Session Law 2022-72, Sec. 9G.7.(a)-(e) or as a covered person under the federal PREP Act functioning as vaccinating providers (collectively “vaccinators”) to administer COVID-19 Vaccines approved or authorized by the FDA per conditions of this order.

Table with 2 columns and 3 main sections: COVID-19 Vaccination, Assessment Criteria, and Plan of Care. Each section contains specific conditions, criteria, and actions for vaccine administration.



**NC State Health Director's Statewide Standing Order
Administration of Moderna COVID-19 Vaccine Bivalent Original and Omicron BA.4 /BA. 5 in Individuals 18
Years of Age and Older
September 2, 2022**

2. Provide Fact Sheet for Recipients and Caregivers for [Moderna COVID-19 Vaccine Bivalent Original and Omicron BA.4 /BA. 5 in Individuals 18 Years of Age and Older](#)
3. [V-safe information sheet](#) to vaccine recipients/caregivers and encourage vaccine recipients to participate in V-safe.

Moderna COVID-19 Vaccine, Bivalent Vaccination Administration Procedures

1. Review Interim [Clinical](#) Considerations for Use of COVID-19 Vaccines Currently Approved or Authorized in the United States
2. [Review CDC EUA for Healthcare Providers of Moderna COVID-19 Vaccine Bivalent Original and Omicron BA.4 /BA. 5 in Individuals 18 Years of Age and Older](#)
3. Appropriate medical treatment and clinical staff able to manage immediate allergic reactions must be immediately available in the event an acute anaphylactic reaction occurs following administration of Moderna Bivalent vaccine.
4. A medical provider, defined as a physician, physician assistant, nurse practitioner, or a pharmacist authorized to order COVID-19 vaccines by the PREP Act must be accessible to provide medical supervision of the vaccination site/service, to assess and evaluate individuals who present with contraindications or precautions to vaccination, and to answer questions or address problems with carrying out this standing order. This may be by telephone or virtual accessibility.
5. Review [Special Circumstances, Precautions, Contraindications, and Criteria or Circumstances for Notifying Medical Provider](#) sections of this standing order **before** administering the bivalent COVID-19 booster vaccine.
6. Following the current [CDC Pre-Vaccination Checklist for COVID-19 Vaccines](#), instruct patients accordingly or consult with overseeing provider.
 - a. The pre-vaccination checklist shall be followed with the following exception: COVID-19 vaccination should **not** be deferred in patients who received monoclonal antibody treatment or convalescent plasma. Patients **should** delay taking EVUSHELD for two weeks after COVID-19 vaccination.
7. Consent must be obtained from the patient prior to vaccine administration per agency policy and in accordance with [NC General Statute. 90-21.13](#).
8. [Personal Protective Equipment](#): Before administering the COVID-19 vaccination, don appropriate personal protective equipment (PPE) per [CDC guidelines for COVID-19 vaccinations](#) to protect against the transmission of COVID-19.




NC State Health Director’s Statewide Standing Order
Administration of Moderna COVID-19 Vaccine Bivalent Original and Omicron BA.4 /BA. 5 in Individuals 18 Years of Age and Older
September 2, 2022

Vaccine Product & Preparation:

1. Vaccine Preparation: Follow the same manufacturer’s guidance for storing/handling vaccine as with previous Moderna vaccine products. Refer to: Moderna COVID-19 Vaccine Preparation and Administration Summary.

2. Presentation:



3. Dosing: 0.5 ml/ 50 mcg/dose

4. Timing:

- a. Timing (interval) between last dose of a COVID-19 primary series shall be at least 2 months.
b. Timing (interval) between last dose of a COVID-19 booster vaccine shall be at least 2 months.

5. Administration:

- a. Route of Administration: Administer vaccine via intramuscular (IM) injection in the deltoid muscle of the arm to patients 18 years of age and older. If contraindications exist to using the deltoid, the anterolateral thigh also can be used.
b. Needle Gauge: Changing needles between drawing up vaccine from a vial and injecting it into a patient is not necessary unless the needle has been damaged, contaminated, or if the needle used to draw up the vaccine is not the correct size for the patient based on their reported weight. Patients may self-report their weight for needle selection purposes.



**NC State Health Director’s Statewide Standing Order
Administration of Moderna COVID-19 Vaccine Bivalent Original and Omicron BA.4 /BA. 5 in Individuals 18
Years of Age and Older
September 2, 2022**

Needle Sizing for Adults			
Sex and Weight of Patient	Needle Gauge	Needle Length	Injection Site*
Female or male fewer than 130 lbs.	22–25	5/8 ** –1"	Deltoid muscle
Female or male 130–152 lbs.	22–25	1"	Deltoid muscle
Female 152–200 lbs.	22–25	1-11/2"	Deltoid muscle
Male 153–260 lbs.	22–25	1-11/2"	Deltoid muscle
Female 200+ lbs.	22–25	11/2"	Deltoid muscle
Male 260+ lbs.	22–25	11/2"	Deltoid muscle

* Alternatively, the anterolateral thigh also can be used.
 ** Some experts recommend a 5/8-inch needle for men and women who weigh less 130 pounds. If used, skin must be stretched tightly (**do not bunch subcutaneous tissue**).

c. **Multiple vaccinations:** If multiple vaccines are administered at a single visit, administer each injection in a different injection site following guidance in the CDC Interim Clinical Considerations on Coadministration of COVID-19 vaccines with other vaccines.

d. **Bleeding Risk:** Patients with blood disorders or who are on blood thinners: administer the vaccine using a 23 gauge or smaller caliber needle, followed by firm pressure on the site, without rubbing, for at least 2 minutes.

6. **Documentation:**

a. **CVMS/NCIR:** Document vaccine record in CVMS or NCIR **within 24 hours** after vaccine administration per system guidelines found at: <https://immunize.nc.gov/providers/covid-19training.htm>. If vaccine is documented in the EHR within 24 hours, providers have **no more than 72 hours** from administration to also enter data in CVMS or NCIR. Determining whether to document in CVMS versus NCIR should occur according to site protocol; it is not necessary to document in both systems.

b. **Electronic Medical Record:** If necessary for billing or other purposes, document patient COVID-19 vaccination in agency electronic medical record per agency policy.

c. Provide vaccine recipients and/or their legal representative COVID-19 Vaccination Record Card indicating the vaccine dose number, product name/manufacturer, lot number, date of vaccination, name/location of vaccinator and clinic site.



NC State Health Director’s Statewide Standing Order
Administration of Moderna COVID-19 Vaccine Bivalent Original and Omicron BA.4 /BA. 5 in Individuals 18 Years of Age and Older
September 2, 2022

Table with 2 columns and 2 rows. Row 1: Moderna COVID-19 Vaccine, Bivalent Observation and Follow-Up. Row 2: Special Circumstances. Content includes observation periods (30 and 15 minutes), anaphylaxis management, and syncopal management.



**NC State Health Director's Statewide Standing Order
Administration of Moderna COVID-19 Vaccine Bivalent Original and Omicron BA.4 /BA. 5 in Individuals 18
Years of Age and Older
September 2, 2022**

	<p>In addition, individuals who received a vaccine that is not listed for emergency use by WHO but for which a U.S. data and safety monitoring board or equivalent has independently confirmed efficacy are considered fully vaccinated.</p> <p>*The following persons require medical consultation.</p> <p>If clinical trial participants have questions about whether they should receive a bivalent booster dose outside of the clinical trial, they should consult with their healthcare provider. Clinical trial participants who did not receive all the recommended doses, or who received other vaccines not listed above, should consult with their healthcare provider to determine if they should receive an FDA-approved or FDA-authorized COVID-19 vaccine series. For more information, refer to Interim Clinical Considerations, Appendix C (People who received COVID-19 vaccine as part of a clinical Trial) outside the United States).</p>
Follow-up	<p>Adverse events that occur in a recipient following COVID-19 vaccination should be reported to VAERS. Vaccination providers are required by the FDA to report the following that occur after COVID-19 vaccination under BLA or EUA:</p> <ul style="list-style-type: none">• Vaccine administration errors<ul style="list-style-type: none">○ Information on preventing, reporting, and managing COVID-19 vaccine administration errors is found in Appendix E. Vaccine administration errors and deviations• Serious adverse events• Cases of Multisystem Inflammatory Syndrome• Cases of COVID-19 that result in hospitalization or death <p>Reporting is encouraged for any other clinically significant adverse event, even if it is uncertain whether the vaccine caused the event. Information on how to submit a report to VAERS is available at https://vaers.hhs.govexternal icon or by calling 1-800-822-7967.</p>
Precautions for Use of this Order	<p>Most people determined to have a precaution to the Moderna COVID-19 Vaccine, Bivalent at their appointment can and should be administered vaccine. See CDC Contraindications and Precautions for COVID-19 Vaccination</p> <ol style="list-style-type: none">1. Persons with a history of an immediate allergic reaction to any other vaccine other than COVID-19 vaccine or to any injectable therapy (i.e., intramuscular, intravenous, or subcutaneous vaccines or therapies [excluding subcutaneous immunotherapy for allergies, i.e., “allergy shots”]). This includes people with a history of an immediate allergic reaction to a vaccine or injectable therapy that contains multiple components, one of which is a COVID-19 vaccine component, even if it is unknown which component elicited the immediate allergic reaction.2. Persons with a contraindication to one type of a COVID-19 vaccine (e.g., viral vector – Janssen/Johnson and Johnson) have a precaution to another (e.g., mRNA – COMIRNATY/Pfizer or Moderna/SPIKEVAX) because of potential cross-reactive hypersensitivity. Consultation with an allergist-immunologist should be considered



**NC State Health Director’s Statewide Standing Order
Administration of Moderna COVID-19 Vaccine Bivalent Original and Omicron BA.4 /BA. 5 in Individuals 18
Years of Age and Older
September 2, 2022**

	<p>prior to vaccination and patients with this precaution should be vaccinated in a health care setting where allergic reactions can be immediately managed and under the supervision of a health care provider experienced in the management of severe allergic reactions.</p> <ol style="list-style-type: none"> 3. Patient self-reported moderate to severe acute illness. 4. Persons with a history of myocarditis or pericarditis after a dose of an mRNA or Novavax COVID-19 vaccine. 5. Persons with a history of MIS-C or MIS-A. 6. Persons with a precaution to vaccination must be counseled about the unknown risks of experiencing a severe allergic reaction and balance these risks against the benefits of vaccination.
<p>Contraindications for Use of this Order</p>	<p>Do not administer the COVID-19 Vaccine to individuals with a history of:</p> <ul style="list-style-type: none"> • Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a component of the vaccine • History of a known diagnosed allergy to a component of the COVID-19 vaccine • Immediate allergic reaction of any severity to a previous dose or known (diagnosed) allergy to a component of the vaccine. <p>See Contraindications and Precautions for COVID-19 Vaccination</p>
<p>Criteria or Circumstances for Notifying Medical Provider</p>	<ol style="list-style-type: none"> 1. Allergic reaction: Call 911, implement medical emergency protocols and immediately notify the medical provider providing clinical supervision of the vaccination site/service. 2. Patient reports a precaution for the vaccine. 3. COVID-19 Vaccine history cannot be determined or is not available. 4. Patients vaccinated with COVID-19 vaccines not authorized or approved in the US. 5. Patients vaccinated with active COVID-19 vaccine as part of a clinical trial. 6. Patient reports they are a HCT or CAR-T cell recipient. These patients may need revaccination, dependent on when the transplant or therapy occurred. 7. Notify the Medical Provider from the organization providing clinical supervision of the vaccination site/service at any time there are questions or problems with carrying out this standing order. <p>Note: Healthcare providers or health departments in the United States can request a consultation from CISA COVID for a complex COVID-19 vaccine safety question that is (1) about an individual patient residing in the United States or vaccine safety issue and (2) not readily addressed by CDC or Advisory Committee on Immunization Practices (ACIP) guidelines.</p>



NC DEPARTMENT OF
**HEALTH AND
HUMAN SERVICES**

**NC State Health Director's Statewide Standing Order
Administration of Moderna COVID-19 Vaccine Bivalent Original and Omicron BA.4 /BA. 5 in Individuals 18
Years of Age and Older
September 2, 2022**

EC Tilson

Approved by: _____
Elizabeth Cuervo Tilson, MD, MPH
NPI: 1760540421

Date Signed: 9-2-22

This standing order is effective immediately and authorizes any North Carolina healthcare provider, in accordance with the conditions of their licensure, or pursuant to orders issued under North Carolina [Session Law 2022-72, Sec. 9G.7.\(a\)-\(e\)](#) or as a covered person under the federal PREP Act functioning as vaccinating providers (collectively "vaccinators") to administer COVID-19 Vaccines authorized by the FDA per conditions of this order.